## American Family: A Syndrome Blood Washer's Syndrome

Blood Blood Blood

There is so much blood on the

walk Blood

Blood

Blood Blood There is so much blood on the walk Blood Blood Blood

> Blood There is so much

on the walk blood

Blood

Blood Blood Blood

There is

much blood SO

> the on

walk Blood

Blbloddllood Blbodb ordisantichbodd blevalk

Blood Blo The

Blood Blood Blood Hereissomuch blood on the walk

Blood Blood Thereissomuch blood on the walk

Blood Blood Thereissomuch blood on the walk

Blood Blood Thereissomuch blood on the walk

Blood Blood Blood Thereissomuch blood on the walk

Blood Blood Blood Blood Thereissomuch blood on the walk

Blood Blood Blood Blood Thereissomuch blood on the walk

Blood There is so much blood on the walk

Blood

Blood Blood Blood There so much blood on the walk Blood is

Temporary Disorder-Triggered by repeated viewings video-recorded police killings, often in a single day, a Blood Washer patient, typically father. develops inexplicable urge to sidewalks. In the early stages of this illness patients spend prolonged moments looking at the floor or their feet. Though early symptoms of the Blood Washer's Syndrome surface in various ways, the disease fully initializes when the patient, in a hypnotic-like trance, wanders outside carrying a bucket. movements become methodical. He will start on any street corner, rinse an area of pavement, apply detergent, rinse the section then continue to the adjacent area and repeat the sequence. During this lapse patients do not pause for food, water, or bathroom breaks. Not to be confused with Coffin Maker's Syndrome, Blood Washers intermittently repeat "Blood. Blood. There is so much Blood on the walk."

Symptoms can exhibit for hours or days. Often times, once the trance lifts subjects unaware that they have been under the Blood Washer's trance. Many feel disoriented and nausea when the trance breaks.

If early signs of the Blood Washer's Syndrome exhibited in a possible patient's behavior it is best to avoid late night television, YouTube, Facebook, and Twitter for two weeks. Ultimately it is best for the patient to submit himself to psychological observation for 48 hours or until early symptoms subside.